

A REVIEW OF THE ESTONIAN DITOMYIIDAE, KEROPLATIDAE, AND DIADOCIDIIDAE (DIPTERA, NEMATOCERA)

Olavi KURINA

Institute of Zoology and Botany, Riia 181, EE-2400 Tartu, Estonia; e-mail: olavi@zbi.ee

Received 16 October 1996, accepted 21 March 1997

Abstract. Data on 19 Estonian species of Ditomyiidae, Keroplatidae, and Diadocidiidae are presented, 8 of them new to the Estonian list of fauna: *Summerus nobilis* Lacksch. (Ditomyiidae); *Cerotelion humeralis* (Zett.), *Monocentrota lundstroemi* Edw., *Orfelia pallida* (Staeg.), *O. unicolor* (Staeg.), *Pyratula zonata* (Zett.) (Keroplatidae); *Diadocidia* (*D.*) *spinosula* Tollet, *D.* (*A.*) *valida* Mik (Diadocidiidae). The northernmost record of *Summerus nobilis* Lacksch. is given.

Key words: Diptera, Ditomyiidae, Keroplatidae, Diadocidiidae, Estonia.

Ditomyiidae, Keroplatidae, and Diadocidiidae are small families among the Nematocera (Diptera). Below these families are dealt in accordance with Mamaev & Krivosheina (1988), Krivosheina & Mamaev (1988), and Krivosheina (1988).

In the Palaearctic region 16 species of Ditomyiidae, 61 species of Keroplatidae, and 5 species of Diadocidiidae have been recorded (Mamaev & Krivosheina, 1988; Krivosheina & Mamaev, 1988; Krivosheina, 1988; Zaitzev, 1994).

In the neighbouring areas of Estonia 2 species of Ditomyiidae (Dit.), 23 species of Keroplatidae (Ker.), and 3 species of Diadocidiidae (Diad.) are known (see the Table). Two species – *Summerus annulatus* (Meigen, 1830) (Dit.) and *Keroplatus tipuloides* Bosc, 1792 (Ker.) – are included in the Red Book of Russian Karelia (Ivanter & Kuznetsov, 1995). *Keroplatus tipuloides* (= *sesidioides* Wahlberg, 1839) is also in Red Lists of Finland and Sweden (Rassi & Väistönen, 1987; Ehnström et al., 1993).

Formerly 11 species of Ditomyiidae, Keroplatidae, and Diadocidiidae were registered in Estonia. According to Krivosheina and Mamaev (1988) *Keroplatus tipuloides* has also been recorded in Estonia, but it is obviously a mistake. In the literature available to me, the species has not been recorded in Estonia.


The species of Ditomyiidae, Keroplatidae, and Diadocidiidae in Estonia and in neighbouring areas

Fin. – Finland, Lat. – Latvia, Len. D., Kar. – Leningrad District and Russian Karelia, Est. – Estonia

Families/Species	Fin.	Lat.	Len. D., Kar.	Est.
	References*			
Ditomyiidae				
<i>Summerus annulatus</i> (Meig.)	1	3	7	3, 10
<i>S. nobilis</i> Lacksch.	–	3	–	10
Keroplatidae				
<i>Asindulum nigrum</i> Latr.	–	3	5	3, 10
<i>Cerotelion humeralis</i> (Zett.)	1	3	5, 6	10
<i>C. lineatus</i> (Fabr.)	1	–	–	–
<i>Isoneuromyia semirufa</i> (Meig.)	1	3	5	3, 10
<i>Keroplatus dispar</i> Dufor	–	–	5	–
<i>K. testaceus</i> (Dalman)	1	3	6, 9	3, 10
<i>K. tipuloides</i> Bosc	1, 2	–	7	–
<i>Macrorrhyncha flava</i> Winn.	1	–	5	4, 10
<i>M. rostrata</i> (Zett.)	1	3	5	–
<i>Monocentrota lundstroemi</i> Edw.	1	–	5	10
<i>Neoplatyura flava</i> (Macq.)	–	3	8	3, 10
<i>N. modesta</i> (Winn.)	1	–	–	–
<i>Orfelia basalis</i> (Winn.)	1	–	–	–
<i>O. discoloria</i> (Meig.)	1	3	5, 9	3
<i>O. fasciata</i> (Meig.)	1	3	–	3, 10
<i>O. nemoralis</i> (Meig.)	1	3	5	3
<i>O. nigricornis</i> (Fabr.)	1	–	5	3
<i>O. pallida</i> (Staeg.)	1	–	–	10
<i>O. unicolor</i> (Staeg.)	1	–	–	10
<i>Platyura bicolor</i> (Macq.)	–	3	–	–
<i>Pyratula zonata</i> (Zett.)	1	–	–	10
<i>Urytalpa ochracea</i> (Meig.)	1	3	–	–
<i>U. trivittata</i> (Lundst.)	1	–	–	–
Diadocidiidae				
<i>Diadocidia (D.) ferruginosa</i> (Meig.)	1	3	5	3, 4, 10
<i>D. (D.) spinosula</i> Tollet	1	–	–	10
<i>D. (A.) valida</i> Mik	1	3	–	10
Total number of species	23	15	15	19

* 1, Hackman, 1980; 2, Ståhls & Kaila, 1990; 3, Lackschewitz, 1937; 4, Dampf, 1924; 5, Zaitzev, 1994; 6, Krivosheina et al., 1986; 7, Ivanter & Kuznetsov, 1995; 8, Yakovlev & Möttus, 1989; 9, Yakovlev, 1986; 10, Original data, and data not published earlier.

The list of Dampf (1924) contains two species, *Diadocidia (D.) ferruginosa* (Meigen, 1830) (Diad.) and *Macrorrhyncha flava* Winnertz, 1846 (Ker.), collected by him from Määvli bog on Hiiumaa Island (see the Figure) and determined by Landrock. Lackschewitz (1937) abstracted Dampf (1924) and gave also data on



Sampling localities. □ data in literature, • original data and data not published earlier. 1, Määvli bog on Hiiumaa Island (Dampf, 1924); 2, Viidumäe Nature Reserve; 3, Islet of Kõinastu near Muhi Island; 4, Piiri on Muhi Island; 5, Suuremõisa on Muhi Island; 6, Klooga, northwest of Keila; 7, Ridala, southeast of Haapsalu (Lackschewitz, 1937); 8, Oonga, southeast of Haapsalu; 9, Kunila, south of Lihula; 10, Puhtu, near Virtsu; 11, Tõstamaa; 12, Audru (Lackschewitz, 1937); 13, Vändra (Lackschewitz, 1937); 14, Jõesuu, northeast of Pärnu; 15, Uulu, south of Pärnu; 16, Rannametsa, south of Pärnu; 17, northeast coast of Lake Rae, southwest of Kilingi-Nõmme; 18, Kabli, south of Pärnu; 19, Nigula Nature Reserve; 20, Ala, west of Tõrva; 21, Tõrva; 22, west coast of Lake Aheru, southeast of Valga; 23, Hargla, southeast of Valga; 24, west coast of Lake Kahrila, southwest of Võru; 25, Hanikase, west of Võru; 26, Kasaritsa, south of Võru (Lackschewitz, 1937); 27, Sangaste; 28, Piigandi, northwest of Võru (Lackschewitz, 1937); 29, Kiuma, southwest of Põlva; 30, Taevaskoja, north of Põlva; 31, Rannaküla, east coast of Lake Võrtsjärv; 32, Vapramäe, northeast of Elva; 33, Kambja, south of Tartu; 34, Järveselja Experimental Forestry Enterprise, southeast of Tartu; 35, Melliste, southeast of Tartu; 36, Tartu and Tartu Tähtvere (Lackschewitz, 1937); 37, Tiksoja, near Tartu; 38, southwest coast of Lake Vasula, near Tartu; 39, Jürküla, southwest of Puurmani; 40, Suuresöödi, southeast of Puurmani; 41, Voore, east of Jõgeva; 42, Endla Nature Reserve; 43, Kose, southeast of Tallinn (Lackschewitz, 1937); 44, Raasiku, east of Tallinn (Lackschewitz, 1937); 45, Revoja, near Palmse in Lahemaa National Park.

ten species collected by Kennel and Sintenis from Ridala, Audru, Vändra, Kasaritsa, Piigandi, Tartu, Tartu Tähtvere, Kose, and Raasiku (Figure). According to Lackschewitz the material of Sintenis included also *Zelmira basalis* Winnertz, 1863 = *Orfelia basalis* (Winnertz, 1863) (Ker.) collected from Audru, and *Zelmira modesta* Winnertz, 1863 = *Neoplatyura modesta* (Winnertz, 1863) (Ker.) collected from Raasiku, but the specimens were later destroyed by pests and they were impossible to check.

In this paper additional material on eight Ditomyiidae, Keroplatidae, and Diadocidiidae species is presented. All Estonian species, except *Summerus nobilis* Lackschewitz, 1937 (Dit.), are widely distributed in Europe (Mamaev & Krivosheina, 1988; Krivosheina & Mamaev, 1988; Krivosheina, 1988; Zaitzev, 1994).

I collected most of the material for the present communication by sweep netting from 31 sites in Estonia (Figure). One specimen of *Cerotelion humeralis* (Zetterstedt, 1850) (Ker.) and one of *Keroplatys testaceus* (Dalman, 1818) (Ker.) were collected by K. Elberg from the inner side of a window in a house at Ala and Hargla. Three specimens of *Diadocidia* (*D.*) *ferruginosa* (Diad.) and one specimen of *D.* (*A.*) *valida* Mik, 1874 (Diad.) were collected by K. Kimmel with a light trap from Endla Nature Reserve. One specimen of *Keroplatys testaceus*, one specimen of *Monocentrota lundstroemi* Edwards, 1925 (Ker.), and two specimens of *Orfelia fasciata* (Meigen, 1804) (Ker.) were collected by H. Remm from Sangaste, Hanikase, and Rannaküla, but the method used is not known.

The material is deposited at the Institute of Zoology and Botany, Tartu, Estonia. Asterisks before the names in the species list indicate new species to Estonia.

LIST OF SPECIES

Ditomyiidae

1. *Summerus annulatus* (Meigen, 1830)

Lackschewitz, 1937: 1 (from Kasaritsa, Audru, Piigandi, and Tartu Tähtvere)

Material: 1 ♂, 25. 06. 1995, Piiri on Muhu Island; 1 ♂, 24. 06. 1996, Melliste. Total 2♂♂.

*2. *S. nobilis* Lackschewitz, 1937

Earlier known from Germany, Latvia, Ukraine, and Belarus (Lackschewitz, 1937; Stackelberg, 1969; Munroe, 1974; Zaitzev, 1994). This record is the northernmost so far.

Material: 1 ♂, 04. 07. 1994, Nigula Nature Reserve.

Keroplatidae

3. *Asindulum nigrum* Latreille, 1805

Lackschewitz, 1937: 7 (from Kasaritsa)

Material: 1♂, 14. 08. 1992, Oonga.

*4. *Cerotelion humeralis* (Zetterstedt, 1850)

Material: 1♂, 18. 06. 1989, Voore; 1♂, 01. 07. 1995, Taevaskoja; 1♂, 14. 07. 1995, Ala, K. Elberg leg. Total 3♂♂.

5. *Isoneuromyia semirufa* (Meigen, 1818)

Lackschewitz, 1937: 7 (as *Zelmira semirufa* Meig. from Ridala)

Material: 3♂♂, 05. 08. 1990, 07. 08. 1990, and 13. 07. 1994, Nigula Nature Reserve.

6. *Keroplatus testaceus* (Dalman, 1818)

Lackschewitz, 1937: 7 (from Tartu, Kasaritsa, Piigandi, and Audru)

Material: 1♂, 05. 07. 1963, Sangaste, H. Remm leg.; 1♀, 17. 08. 1991, Puhtu; 1♂, 09. 07. 1994, Hargla, K. Elberg leg.; 1♂, 25. 08. 1995, Tõrva. Total 3♂♂ 1♀.

7. *Macrorrhyncha flava* Winnertz, 1846

Dampf, 1924: 44 (as *Asindulum flavum* Winn. from Määvli bog on Hiiumaa Island)

Material: 1♂, 21. 08. 1991, Nigula Nature Reserve.

*8. *Monocentrota lundstroemi* Edwards, 1925

Material: 1♂, 25. 07. 1985, Hanikase, H. Remm leg.; 1♀, 26. 06. 1996, Jõesuu. Total 1♂ 1♀.

9. *Neoplatyura flava* (Macquart, 1826)

Lackschewitz, 1937: 7 (as *Zelmira flava* Macq. from Audru and Ridala)

Material: 19♂♂ 4♀♀, Piiri on Muhu Island (1995), Oonga (1992, 1995), Nigula Nature Reserve (1990, 1993, 1995, 1996), Kiuma (1995), Tiksoja (1994).

10. *Orfelia discoloria* (Meigen, 1818)

Lackschewitz, 1937: 8 (as *Zelmira discoloria* Meig. from Tartu, Kasaritsa, Audru, Raasiku).

11. *O. fasciata* (Meigen, 1804)

Lackschewitz, 1937: 7 (as *Zelmira fasciata* Meig. from Tartu, Audru, Kose)

Material: 1♂, 26. 06. 1956, Rannaküla, H. Remm leg.; 1♀, 13. 07. 1957, Sangaste, H. Remm leg.; 1♂, 17. 06. 1989, Voore; 1♂, 01. 07. 1996, Suuremõisa on Muhu Island. Total 3♂♂ 1♀.

12. *O. nemoralis* (Meigen, 1818)

Lackschewitz, 1937: 8 (as *Zelmira nemoralis* Meig. from Tartu)

13. *O. nigricornis* (Fabricius, 1805)

Lackschewitz, 1937: 7 (as *Zelmira nigricornis* Fabr. from Audru and Vändra)

*14. *O. pallida* (Staeger, 1840)

Material: 1♂, 03. 07. 1994, Nigula Nature Reserve.

*15. *O. unicolor* (Staeger, 1840)

Material: 1♂ 1♀, 04. 07. 1994 and 11. 07. 1993, Nigula Nature Reserve; 1♂, 18. 06. 1989, Voore. Total 2♂♂ 1♀.

*16. *Pyratula zonata* (Zetterstedt, 1855)

Material: 1♂, 26. 06. 1993, Viidumäe Nature Reserve; 1♂, 12. 06. 1994, Taevaskoja; 1♀, 02. 07. 1994, Nigula Nature Reserve; 1♀, 03. 07. 1994, coast of Lake Rae. Total 2♂♂ 2♀♀.

Diadocidiidae

17. *Diadocidia (Diadocidia) ferruginosa* (Meigen, 1830)

Dampf, 1924: 44 (from Määvli bog on Hiiumaa Island)

Lackschewitz, 1937: 6 (from Tartu, Audru, and Kose)

Material: 42♂♂ 3♀♀, Viidumäe Nature Reserve (1988), Kõinastu (1994), Oonga (1993), Klooga (1996), NE coast of Lake Rae (1994), Nigula Nature Reserve (1990, 1991, 1992, 1993, 1994, 1995), Hargla (1994), W coast of Lake Aheru (1994), W coast of Lake Kahrila (1995), Vapramäe (1995), Kambja (1995), Järvselja (1989), Melliste (1994), SW coast of Lake Vasula (1995), Jüriküla (1995), Suuresöödi (1994), Voore (1989); 3♂♂, 04.–11. 06. 1995, Endla Nature Reserve, K. Kimmel leg. Total 45♂♂ 3♀♀.

*18. *D. (D.) spinosula* Tollet, 1948

Material: 17♂♂ 7♀♀, Piiri on Muhu Island (1995, 1996), Oonga (1989, 1993), Klooga (1996), Kunila (1995), Tõstamaa (1994), Uulu (1995), Rannametsa (1995), Kabli (1995), Nigula Nature Reserve (1991, 1994, 1995), Hargla (1994), Vapramäe (1989, 1995), Lahemaa, Revoja (1996).

*19. *D. (Adidocida) valida* Mik, 1874

Material: 1♂, 04. 07. 1994, Nigula Nature Reserve; 1♂, 8.–15. 10. 1995, Endla Nature Reserve, K. Kimmel leg. Total 2♂♂.

ACKNOWLEDGEMENTS

I express my thanks to the Estonian Science Foundation for financial support, grant No. 128. I am greatly indebted to Mart Jüssi and Robert Oetjen for reading the manuscript.

REFERENCES

- Dampf, A. 1924. Zur Kenntnis der estländischen Hochmoorfauna, I. *Beiträge zur Kunde Estlands*. Reval, 10, 33–49.
- Ehnström, B., Gärdenfors, U. & Lindelöw, Å. 1993. *Rödlistade evertebrater i Sverige 1993*. Databanken för hotade arter, Uppsala.
- Hackman, W. 1980. A check list of the Finnish Diptera I. Nematocera and Brachycera (s. str.). *Notulae Entomologicae*, **60**, 17–48.
- Ivanter, E. V. & Kuznetsov, O. L. 1995. *Red Book of Karelia*. Karelia, Petrozavodsk (in Russian).
- Krivosheina, N. P. 1988. Family Diadocidiidae. In *Catalogue of Palaearctic Diptera*. Vol. 3. *Ceratopogonidae–Mycetophilidae* (Soos, Å. & Papp, L., eds.). Akadémiai Kiadó, Budapest, 210–211.
- Krivosheina, N. P. & Mamaev, B. M. 1988. Family Keroplatidae. In *Catalogue of Palaearctic Diptera*. Vol. 3. *Ceratopogonidae – Mycetophilidae* (Soos, Å. & Papp, L., eds.). Akadémiai Kiadó, Budapest, 199–210.
- Krivosheina, N. P., Zaitzev, A. I. & Yakovlev, E. B. 1986. *Insects as Decomposers of Fungi in the Forest of the European Part of the USSR*. Nauka, Moscow (in Russian).
- Lackschewitz, P. 1937. Die Fungivoridae des ostbaltischen Gebites. *Arb. Naturf. – Ver. Riga*, N. F. 21, 1–47.
- Mamaev, B. M. & Krivosheina, N. P. 1988. Family Ditomyiidae. In *Catalogue of Palaearctic Diptera*. Vol. 3. *Ceratopogonidae – Mycetophilidae* (Soos, Å. & Papp, L., eds.). Akadémiai Kiadó, Budapest, 197–199.
- Munroe, D. D. 1974. The systematics, phylogeny, and zoogeography of *Symmerus* Walker and *Australosymmerus* Freeman (Diptera: Mycetophilidae: Ditomyiinae). *Memoirs of the Entomological Society of Canada*, **92**.
- Rassi, P. & Väisänen, R. 1987. *Threatened Animals and Plants in Finland*. Valtion painatuskeskus, Helsinki.
- Stackelberg, A. A. 1969. Fam. Ditomyiidae. In *Key to the Insects of the European Part of the USSR*. V. Part I. (Bei-Bienko, G. Ya., ed.). Nauka, Leningrad (in Russian).
- Ståhls, G. & Kaila, L. 1990. *Keroplatys tipuloides* Bosc rediscovered in Finland (Diptera: Nematocera: Keroplatidae). *Notulae Entomologicae*, **69**, 203–206.
- Yakovlev, E. B. 1986. Fungivorous insects of South Karelia (ecological and faunal list). In *Fauna and Ecology of Karelian Arthropods*. Institute of Forest, Petrozavodsk (in Russian).
- Yakovlev, E. B. & Möttus, E. P. 1989. *On the Attraction of Insects with Fruit Bodies of Mushrooms and by Different Components of Mushrooms Scent*. Institute of Forest, Petrozavodsk (in Russian).
- Zaitzev, A. I. 1994. *Fungus Gnats of the Fauna of Russia and Adjacent Regions*. Part 1. Nauka, Moscow (in Russian).

ÜLEVAADE SUGUKONDADEST DITOMYIIDAE, KEROPLATIDAE JA DIADOCIDIIDAE (DIPTERA, NEMATOCERA) EESTIS

Olavi KURINA

On esitatud andmed sugukondade Ditomyiidae, Keroplatidae ja Diadocidiidae Eestis esineva 19 liigi kohta. Nendest kahekse on leitud Eestis esmakordelt: *Summerus nobilis* Lacksch. (Ditomyiidae); *Cerotelion humeralis* (Zett.), *Monocentrota lundstroemi* Edw., *Orfelia pallida* (Staeg.), *O. unicolor* (Staeg.), *Pyratula zonata* (Zett.) (Keroplatidae); *Diadocidia* (*D.*) *spinosa* Tollet, *D. (A.) valida* Mik (Diadocidiidae). Liigi *Summerus nobilis* Lacksch. leid on siiani kõige põhjapoolsem.

TABLE I. ESTIMATE OF THE TOTAL AND AVERAGE ANNUAL CATCHES OF FISH IN LAKE PEIPUS

Catches of fish	Year	Total catch, 10 ³ kg	Average annual catch, 10 ³ kg	Relative abundance of fish	Percentage of fish to total weight of fish	Percentage of fish to total weight of fish and its price	Percentage of fish to total weight of fish and its price				
					1980	1981	1982	1983	1984	1985	1986
Total	1980	557,1	92,8	2.1	100.0	100.0	100.0	100.0	100.0	100.0	100.0
Mus	1980	462,5	77,1	1.0	83.0	83.0	83.0	83.0	83.0	83.0	83.0
Pike	1980	12,4	2,1	0.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
Trout	1980	10,3	1,8	0.2	1.8	1.8	1.8	1.8	1.8	1.8	1.8
Perch	1980	5,2	0,9	0.1	0.9	0.9	0.9	0.9	0.9	0.9	0.9
Chub	1980	3,7	0,6	0.1	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Whitefish	1980	3,5	0,6	0.1	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Angler	1980	2,6	0,4	0.1	0.4	0.4	0.4	0.4	0.4	0.4	0.4
Other fish	1980	20,9	3,5	0.3	3.6	3.6	3.6	3.6	3.6	3.6	3.6
Total	1981	540,8	90,1	2.1	98.2	98.2	98.2	98.2	98.2	98.2	98.2
Mus	1981	449,6	74,9	1.0	83.0	83.0	83.0	83.0	83.0	83.0	83.0
Pike	1981	13,7	2,3	0.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
Trout	1981	11,1	1,8	0.2	1.6	1.6	1.6	1.6	1.6	1.6	1.6
Perch	1981	5,9	0,9	0.1	0.8	0.8	0.8	0.8	0.8	0.8	0.8
Chub	1981	4,1	0,6	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Whitefish	1981	3,9	0,6	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Angler	1981	2,7	0,4	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Other fish	1981	22,4	3,7	0.3	3.6	3.6	3.6	3.6	3.6	3.6	3.6
Total	1982	528,5	88,0	2.1	96.4	96.4	96.4	96.4	96.4	96.4	96.4
Mus	1982	446,3	74,4	1.0	83.0	83.0	83.0	83.0	83.0	83.0	83.0
Pike	1982	14,6	2,4	0.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
Trout	1982	12,0	2,0	0.2	1.8	1.8	1.8	1.8	1.8	1.8	1.8
Perch	1982	6,3	1,1	0.1	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Chub	1982	4,4	0,7	0.1	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Whitefish	1982	3,7	0,6	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Angler	1982	2,5	0,4	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Other fish	1982	21,4	3,5	0.3	3.4	3.4	3.4	3.4	3.4	3.4	3.4
Total	1983	512,5	85,4	2.1	93.6	93.6	93.6	93.6	93.6	93.6	93.6
Mus	1983	434,3	72,3	1.0	83.0	83.0	83.0	83.0	83.0	83.0	83.0
Pike	1983	15,7	2,6	0.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
Trout	1983	10,4	1,7	0.2	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Perch	6,6	1,1	0.1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Chub	4,6	0,7	0.1	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Whitefish	3,7	0,6	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Angler	2,4	0,4	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Other fish	1983	20,7	3,4	0.3	3.3	3.3	3.3	3.3	3.3	3.3	3.3
Total	1984	500,2	83,3	2.1	91.2	91.2	91.2	91.2	91.2	91.2	91.2
Mus	1984	427,0	71,1	1.0	83.0	83.0	83.0	83.0	83.0	83.0	83.0
Pike	1984	16,6	2,7	0.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
Trout	1984	9,6	1,6	0.2	1.4	1.4	1.4	1.4	1.4	1.4	1.4
Perch	6,6	1,1	0.1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Chub	4,6	0,7	0.1	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Whitefish	3,7	0,6	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Angler	2,3	0,4	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Other fish	1984	20,1	3,3	0.3	3.2	3.2	3.2	3.2	3.2	3.2	3.2
Total	1985	485,1	81,2	2.1	88.9	88.9	88.9	88.9	88.9	88.9	88.9
Mus	1985	421,9	70,3	1.0	83.0	83.0	83.0	83.0	83.0	83.0	83.0
Pike	1985	17,4	2,8	0.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
Trout	1985	9,8	1,6	0.2	1.4	1.4	1.4	1.4	1.4	1.4	1.4
Perch	6,8	1,1	0.1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Chub	4,8	0,7	0.1	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Whitefish	3,7	0,6	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Angler	2,3	0,4	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Other fish	1985	19,8	3,2	0.3	3.1	3.1	3.1	3.1	3.1	3.1	3.1
Total	1986	468,9	79,8	2.1	87.2	87.2	87.2	87.2	87.2	87.2	87.2
Mus	1986	418,7	70,1	1.0	83.0	83.0	83.0	83.0	83.0	83.0	83.0
Pike	1986	18,1	2,9	0.2	2.2	2.2	2.2	2.2	2.2	2.2	2.2
Trout	1986	10,3	1,7	0.2	1.5	1.5	1.5	1.5	1.5	1.5	1.5
Perch	6,8	1,1	0.1	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0
Chub	4,8	0,7	0.1	0.6	0.6	0.6	0.6	0.6	0.6	0.6	0.6
Whitefish	3,7	0,6	0.1	0.5	0.5	0.5	0.5	0.5	0.5	0.5	0.5
Angler	2,3	0,4	0.1	0.3	0.3	0.3	0.3	0.3	0.3	0.3	0.3
Other fish	1986	19,1	3,1	0.3	3.0	3.0	3.0	3.0	3.0	3.0	3.0

LAKE PEIPUS

Earlier, L. Peipsi was regarded as a mixed-freedom lake, since the second half of 1980s 2.2-2.5 kg wallers were taken (Fig. 1-7). In the first half of the 1980s, however, the catches of perch and roach were very low (Fig. 1-7), whereas the catches of chub, whitefish and pike increased (Fig. 1-7). The catches of mus and trout were stable (Fig. 1-7). The catches of chub and whitefish decreased in 1984-1985 (Fig. 1-7), but increased again in 1986 (Fig. 1-7). The catches of chub and whitefish were the lowest in 1986 (Fig. 1-7), but the catches of mus and trout were the highest in 1986 (Fig. 1-7). The catches of chub, whitefish and mus were highest in 1980-1983 (Fig. 1-7). The catches of chub, whitefish and trout were the lowest in 1980-1983 (Fig. 1-7).